

Tips to aid oral feeding in children with Chronic Kidney Disease (CKD)

Children with CKD struggle with eating and drinking due to a variety of reasons associated with the condition including having a poor appetite, suffering from nausea, vomiting or acid reflux. These children can often miss key developmental feeding experiences which can make feeding and drinking a challenge. The below points are tips which can be discussed with parents to encourage oral diet for these children.

1. Ensure realistic expectations are discussed with the family to help reduce anxiety and build a positive feeding environment. Encourage and praise little steps; playing with food and having contact with food is a positive step.



2. Encourage families to prepare food together. This will create a positive relationship with food for the child.



3. Encourage the family to all eat together so the child experiences all the sights, smells and sounds of mealtimes¹. Observing others eating and eating with peers can encourage a child to have familiar foods².
4. It is important to keep food experiences positive and enjoyable. Advise parents to stop feeding the child at a mealtime if they become upset or distressed.
5. Allow the child to play with foods in their own time, without pressure. The aim is to create positive eating and drinking experiences.



6. Remove distractions at mealtimes; for example turn off the television, remove ipads and phones³.
7. Limit mealtimes to 20-30 minutes⁴ and avoid too much focus on food.
8. Have snacks available during the day to encourage eating⁵.
9. Ensure the dietary advice provided highlights foods which the child can have as well as those foods which need to be restricted.



Innovation in Nutrition

A Nestlé Health Science Company

© Reg. Trademarks of Société des Produits Nestlé S.A.
© Société des Produits 2018. All rights reserved.

Vitaflor International Ltd,
Suite 1.11,
South Harrington Building,
182 Sefton Street,
Brunswick Business Park,
Liverpool, L3 4BQ, UK.

Nutritional Helpline: **+44 (0) 151 702 4937**

www.vitafloweb.com

All information correct at the time of print

**This information sheet is only intended for
Healthcare Professionals**

References

1. Krom H, de Winter JP, Kindermann A. Development, prevention, and treatment of feeding tube dependency. *European journal of pediatrics*. 2017;176(6):683-8.
2. Mason SJ, Harris G, Blissett J. Tube Feeding in Infancy: Implications for the Development of Normal Eating and Drinking Skills. *Dysphagia*. 2005;20(1):46-61.
3. Infokid. Feeding your baby or child [Available from: <http://www.infokid.org.uk/feeding-your-baby-or-child-chronic-kidney-disease>.

4. Smith. Chapter 28 Faltering weight. In: Shaw, editor. *Clinical Paediatric Dietetics*. 4th: Wiley Blackwell; 2015. p. 764-73.

5. Strologo LD, Principato F, Sinibaldi D, Appiani AC, Terzi F, Dartois AM, et al. Feeding dysfunction in infants with severe chronic renal failure after long-term nasogastric tube feeding. *Pediatric Nephrology*. 1997;11(1):84-6.